Zootaxa 3023: 68 (2011)
www.mapress.com/zootaxa/
Copyright © 2011 · Magnolia Press

Erratum

ISSN 1175-5326 (print edition)
ZOOTAXA
ISSN 1175-5334 (online edition)

Sandoval, C.M., Nieves, E., Angulo, V.M., Rosa, J.A. & Aldana, E. (2011) Morphology of the eggs of the genus *Belminus* (Hemiptera: Reduviidae: Triatominae) by optical and scanning electron microscopy. *Zootaxa*, 2970, 33–40.

On page 33.
The average size of the egg among these species is 1.03–1.17 mm in length and 0.72–0.8 mm in width. [Not 0.51– 0.58 mm in length and 0.36–0.4 mm in width].

On page 35.
For Table 1 the correct values in millimeters of measures of length and width of eggs of three species are: *B. ferroae* (L=1.17±0.07; W=0.80 ±0.03), *B. corredori* (L=1.06±0.04; W=0.77±0.03), *B. herreri* (L= 1.03±0.03; 0.72±0.02). [Not *B. ferroae* (L=0.59±0.03; W=0.40 ±0.01), *B. corredori* (L=0.53±0.02; W=0.39±0.02), *B.herreri* (L= 0.52±0.02; 0.36±0.01)]. L=length, W= width, ±SD= Standard Deviation.